

Correction to “Radiance and flux simulations for mineral dust aerosols: Assessing the error due to using spherical or spheroidal model particles”

M. Kahnert and A. Kylling

Norwegian Institute for Air Research (NILU), Kjeller, Norway

Received 3 August 2004; accepted 11 August 2004; published 19 October 2004.

INDEX TERMS: 9900 Corrections; 0305 Atmospheric Composition and Structure: Aerosols and particles (0345, 4801); 0360 Atmospheric Composition and Structure: Transmission and scattering of radiation; 0669 Electromagnetics: Scattering and diffraction; 3359 Meteorology and Atmospheric Dynamics: Radiative processes; *KEYWORDS:* mineral dust, radiative forcing, remote sensing

Citation: Kahnert, M., and A. Kylling (2004), Correction to “Radiance and flux simulations for mineral dust aerosols: Assessing the error due to using spherical or spheroidal model particles,” *J. Geophys. Res.*, *109*, D20203, doi:10.1029/2004JD005311.

[1] In the paper “Radiance and flux simulations for mineral dust aerosols: Assessing the error due to using spherical or spheroidal model particles” (*Journal of Geophysical Research*, *109*, D09203, doi:10.1029/2003JD004318), all incidences of spectral radiance units $\text{W m}^{-2} \text{nm}^{-1} \text{sr}^{-1}$ should, of course, be replaced by $\text{mW m}^{-2} \text{nm}^{-1} \text{sr}^{-1}$. Likewise, all incidences of spectral

flux units of $\text{W m}^{-2} \text{nm}^{-1}$ have to be replaced by $\text{mW m}^{-2} \text{nm}^{-1}$.

M. Kahnert and A. Kylling, Norwegian Institute for Air Research (NILU), P.O. Box 100, 2027 Kjeller, Norway. (michael.kahnert@nilu.no; arve.kylling@nilu.no)